

BREMELANOTIDE/OXYTOCIN

Any medication questions? Pharmacy counseling available at 866-206-7936.

Bremelanotide (PT-141): PT-141, also known as bremelanotide, is a synthetic peptide that is used to treat sexual dysfunction in men and women. While Viagra and Cialis, which are PDE-5 inhibitors that work to increase blood flow into the penis, PT-141 is a melanocortin, which works in the brain to increase sexual desire. The synthetic peptide blend was developed from Melanotan II, and was originally intended to darken skin pigmentation and induce tanning. However, during the initial testing, patients noted a pleasant side effect - that is caused increased sexual arousal. Sometimes, for men with erectile dysfunction or other sexual performance disorders, a combination of Viagra or Cialis and PT-141 can be effective.

Oxytocin: Oxytocin is a nano peptide hormone that is secreted by the posterior pituitary. Oxytocin produces a wide range of effects both peripherally and locally in the brain. It is released in large amounts in females during childbirth, nipple stimulation, and breastfeeding. Oxytocin is released by males and females during skin-to-skin contact, sexual arousal, and orgasm/ejaculation. Oxytocin causes dilation of vascular smooth muscle, thus increasing renal, coronary, and cerebral blood flow. Blood pressure usually remains unaffected, but with the administration of very large doses or high concentration solutions, blood pressure may decrease transiently. This transient decrease in blood pressure leads to reflex tachycardia and an increase in cardiac output; any fall in blood pressure is usually followed by a small, but sustained, increase in blood pressure. Oxytocin improves sexual enjoyment. Oxytocin will help a man with his ejaculation and orgasm. Other benefits of oxytocin and oxytocin as a libido treatment: causes feelings of intimacy and closeness, triggers increased orgasm intensity, shortens time to orgasm, makes it easier to climax, increases production of vaginal secretions, helps with pleasurable sensations, aids feeling of well-being, warmth, trust and contentment, aids in social anxiety and sociability, helps increase bonding, nurturing and trust.

HOW TO TAKE THIS MEDICATION: Allow the troche to dissolve under your tongue or between the cheek and gum. Do not eat or drink while the medication is dissolving to allow for maximum absorption through the oral mucosa.

SIDE EFFECTS: Dizziness, headache, flushing, or stomach upset may occur. Vision changes such as increased sensitivity to light, blurred vision, or telling blue and green colors apart may also occur. If any of these effects persist or worsen, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly from a sitting or lying position. Sexual activity may put an extra strain on your heart, especially if you have heart problems. If you have heart problems and experience any of these side effects while having sex, stop and get medical help right away: severe dizziness, fainting, chest/jaw/left arm pain, nausea.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. Bremelanotide can cause nausea, which may be severe. Use this medication exactly as your doctor has prescribed.

Call your doctor immediately if you have: severe or ongoing nausea; slow heartbeats; or high blood pressure - severe headache, blurred vision, pounding in your neck or ears.

PRECAUTIONS: Before using this medication, tell your doctor or pharmacist your medical history, especially of: heart problems (such as heart attack or life-threatening irregular heartbeat in the past 6 months, chest pain/angina, heart failure), stroke in the past 6 months, kidney disease, liver disease, high or low blood pressure, dehydration, penis conditions (such as angulation, fibrosis/scarring, Peyronie's disease), history of painful/prolonged erection (priapism), conditions that may increase the



TOLL FREE 866-206-7936
scripts@atlasrxpharmacy.com

risk of priapism (sickle cell anemia, leukemia, multiple myeloma), eye problems (retinitis pigmentosa, sudden decreased vision, NAION), bleeding disorders, active stomach ulcers.

This drug may make you dizzy or cause vision problems. ALCOHOL OR MARIJUANA (cannabis) may make you more dizzy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. LIMIT ALCOHOLIC BEVERAGES. Talk to your doctor if you are using marijuana (cannabis).

You should not use bremelanotide if you have heart problems or if you have untreated or uncontrolled high blood pressure. Tell your doctor if you have ever had: Heart problems; high blood pressure; kidney disease; or liver disease.

DRUG INTERACTIONS: A product that may interact with this drug is: Ephedra, Ma Huang
If you are also taking blood pressure medications to treat high blood pressure, your blood pressure may get too low which can lead to dizziness or fainting.

STORAGE: Refrigerate or store at controlled room temperature and away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets.