

Eros Max Dose

Sildenafil 110 mg / Tadalafil 22 mg / Apomorphine 3 mg

This medication is a **maximum-strength combination treatment** for erectile dysfunction (ED). It contains three active ingredients that work together to improve erections. Because this is a very strong formula, it is extremely important to follow all safety instructions carefully.

What This Medication Contains

Sildenafil 110 mg (*Same active ingredient as Viagra*)

- Helps increase blood flow to the penis
- Works best when taken on an empty stomach
- Requires sexual stimulation to work

Tadalafil 22 mg (*Same active ingredient as Cialis*)

- Increases blood flow to help get and keep an erection
- Lasts longer in the body than sildenafil
- May also help improve urinary symptoms from an enlarged prostate (BPH), such as weak stream or frequent urination

Apomorphine 3 mg

- Works in the brain to stimulate sexual desire and trigger the erection response
- Helps improve arousal
- May increase nausea more than the other ingredients

VERY IMPORTANT: Do NOT Take With Nitrates

Do **NOT** use this medication if you take nitrates for chest pain or heart problems, including:

- Nitroglycerin
- Isosorbide mononitrate
- Isosorbide dinitrate

Combining these can cause a sudden and life-threatening drop in blood pressure.

How to Take This Medication

- Take one dose 30–60 minutes before sexual activity
- Do **NOT** take more than one dose in 24 hours
- Allow the tablet/melt to dissolve under your tongue or between your cheek and gum
- Do not eat or drink while it is dissolving
- Do not chew or swallow whole
- Sexual stimulation is required for it to work
- A heavy or fatty meal may delay how quickly it starts working

Possible Side Effects

Because this is a **maximum-dose combination**, side effects may be more pronounced.

Common side effects may include:

- Headache
- Flushing (warmth/redness of face or neck)
- Dizziness or lightheadedness
- Nausea or upset stomach
- Stuffy nose
- Sensitivity to light
- Blurred vision or blue/green color changes

To reduce dizziness, stand up slowly from sitting or lying down.

Limit alcohol, as it can increase dizziness, fainting, and fast heartbeat.



Seek Emergency Medical Help Immediately If You Have:

- An erection lasting longer than 4 hours (can cause permanent damage)
- Sudden vision loss in one or both eyes
- Sudden hearing loss or ringing in the ears
- Chest pain, jaw pain, left arm pain, shortness of breath, fainting, or severe dizziness during or after sex

Stop taking the medication and get medical help right away if any of these occur.

Important Precautions

Tell your doctor if you:

- Have a history of heart attack, stroke, or irregular heartbeat
- Have high blood pressure or take blood pressure medications
- Take alpha-blockers for prostate problems
- Have diabetes or high cholesterol
- Are over age 50 and smoke
- Have been told you have “crowded disc” or other eye problems

Sexual activity can put extra strain on the heart. If you have heart problems, talk to your doctor before using this medication.

Questions?

Pharmacy counseling is available at **866-206-7936**.

If you are unsure whether this medication is safe for you, speak with your doctor or pharmacist before taking it.